



Contribution ID: 155 Contribution code: S04-AMP-201

Type: Poster presentation

## Nonrelativistic energy spectra of three-particle systems calculated in the adiabatic approach by using finite difference method

*Tuesday, 30 August 2022 18:00 (1h 30m)*

Precise spectroscopy of light atomic and molecular systems gives us new insights in the fundamental laws and physical constants of nature [1]. In particular, exotic atoms like antihydrogen and antiprotonic helium could also allow comparison between matter and antimatter characteristics and test of CPT symmetry [2]. From theoretical point of view, development of very accurate techniques like adiabatic and variational methods give us the tools to compute different atomic quantities with high precision [3].

Here, we present accurate calculations of nonrelativistic energy levels of three-particle systems, like for example  $\bar{p}\text{He}^+$ , in adiabatic approximation. In our approach, the electronic wavefunctions are computed by using two-dimensional finite difference method (FDM) and the radial wavefunctions are obtained by solving a system of eigenvalue problems [4] by iterative application of one-dimensional FDM. We analyze the results of using the method with Cartesian and logarithmic stencils and different number of node points. We show that the level of precision allows our results to be used in computations of leading relativistic corrections to transition spectra of these systems and accurately determining the effects of interactions with external fields. A comparison with other computations of light atomic systems' energy spectra has been made and the advantages and disadvantages have been assessed.

**Acknowledgement:** This work was supported by the Bulgarian Science Fund under contract KP-06-M58/3 / 22.11.2021.

### References

1. I.V. Kortunov et. al., Proton–electron mass ratio by high-resolution optical spectroscopy of ion ensembles in the resolved-carrier regime, *Nature Physics* 17, 569–573 (2021).
2. The ALPHA Collaboration, Investigation of the fine structure of antihydrogen, *Nature*, 578, 375–380 (2020).
3. P. Danev, D. Bakalov, V.I. Korobov, S. Schiller, Hyperfine structure and electric quadrupole transitions in the deuterium molecular ion, *Phys. Rev. A* 103 (1), 012805 (2021).
4. S.I. Vinitskij, L.I. Ponomarev, Adiabatic representation of three-body problem with the Coulomb interaction, *J. Phys. B.* (1978).

**Primary authors:** BAKALOV, Dimitar (Institute for Nuclear Research and Nuclear Energy, Bulgarian Academy of Sciences); TONCHEV, Hristo (Institute of Solid State Physics - Bulgarian Academy of Sciences); DANEV, Petar (Institute for Nuclear Research and Nuclear Energy, Bulgarian Academy of Sciences); STOILOV, Michail (Institute for Nuclear Research and Nuclear Energy, Bulgarian Academy of Sciences)

**Presenter:** DANEV, Petar (Institute for Nuclear Research and Nuclear Energy, Bulgarian Academy of Sciences)

**Session Classification:** Poster session

**Track Classification:** Scientific Sections: S04 Atomic and Molecular Physics