



Contribution ID: 216 Contribution code: S06-CMPSP-221

Type: Poster presentation

Bose-Einstein condensate - Tunneling

Tuesday, August 30, 2022 6:00 PM (1h 30m)

Bose-Einstein condensate is state of matter that is the future of the quantum computers. One of the crucial properties of Bose-Einstein condensate is quantum tunneling. In this paper are listed previously achieved results by other authors considering the factors that affect tunneling in Bose-Einstein condensate, such as form of interaction between particles, shape of potential well and the form and the dimension of the system [1]. According to Meng *et al.* who studied two-dimensional honeycomb optical lattice, tunneling occurs in an attractive interaction regime if dipole gap solitons are in-phase and in repulsive interaction regime if dipole gap solitons are out of phase [2]. If the particle experiences gradual change of potential barrier the probability of the excitation is lower [3]. If the particle experiences sudden change of the potential barrier, probability increases. ³ The further study of the factors that affect tunneling in Bose-Einstein condensate may be conducted in exploring different systems and mixtures of aforementioned condensates.

References

1. X. Zhao *et al.*, Phys. Rev. A **96**, (2017).
2. H. Meng *et al.*, Physica A: Stat. Mech. Appl. **577**, 126087 (2021).
3. D. Lindberg *et al.*, arXiv:2110.15298v3 [cond-mat.quant-gas]

Primary author: PEJIĆ, Emilija (University of Novi Sad, Faculty of Sciences, Department of Physics)

Presenter: PEJIĆ, Emilija (University of Novi Sad, Faculty of Sciences, Department of Physics)

Session Classification: Poster session

Track Classification: Scientific Sections: S06 Condensed Matter Physics and Statistical Physics