




# Measurement of the Higgs to $\gamma\gamma$ branching fraction at 3 TeV CLIC



Goran Kačarević, Ivanka Božović, Nataša Vukašinić  
Vinca Institute of Nuclear Sciences, University of Belgrade, Serbia

## 1. Introduction



- Higgs boson couples with photons on a loop level – this channel is sensitive to BSM contributions
- Higgs decay to two photons is a rare process:  $BR(H \rightarrow \gamma\gamma) \sim 0.23\%$  implying low signal rate and challenging signal separation.



- Typical  $g_{H\gamma\gamma}$  deviations in the BSM models of the Higgs sector are  $\leq 4\%$
- Utmost statistical precision of  $g_{H\gamma\gamma}$  is obtained in a global fit (model independent, k-framework, EFT) is  $\sim 1\%$ .

## 2. CLIC Detector

- 4 T super-conducting solenoid
- Detector system for track reconstruction is based on Si technology
- **Reconstruction and identification of particles using Particle Flow Algorithm (PFA)**
- Excellent performance of photon reconstruction and identification due to the highly granular ECAL:
  - Photon identification efficiency  $\sim 99\%$
  - Photon energy resolution is  $2\% - 3\%$




Photon identification efficiency

Photon energy resolution as a function of number of ECAL layers

## 3. Signal separation

- WW-fusion is dominant Higgs production mechanism for energies above 500 GeV, and provides access to rare Higgs decays ( $\gamma\gamma$ ,  $Z\gamma$ ,  $\mu\mu$ )




- $\sigma(H\nu\nu)$  at 3 TeV is 415 fb
- $\sigma(H\nu\nu) \times BR(H \rightarrow \gamma\gamma) = 0.95$  fb
- $N_{\text{signal}} \sim 4750 \text{ evt}/5ab^{-1}$

| Process   | $\sigma_{\text{effective}}$ (fb) | No. evt, $5 ab^{-1}$ |
|---|----------------------------------|----------------------|
| $\sigma(h\nu\nu) \times BR(h \rightarrow \gamma\gamma)$ | 0.95                             | 4750                 |
| $e^+e^- \rightarrow \gamma\gamma$                       | 15.2                             | $7.6 \cdot 10^4$     |
| $e^+e^- \rightarrow e^+e^-\gamma$                       | 335                              | $1.7 \cdot 10^6$     |
| $e^+e^- \rightarrow e^+e^-\gamma\gamma$                 | 33                               | $1.5 \cdot 10^5$     |
| $e^+e^- \rightarrow \nu\nu\gamma$                       | 13                               | $6.5 \cdot 10^4$     |
| $e^+e^- \rightarrow \nu\nu\gamma\gamma$                 | 26                               | $1.3 \cdot 10^5$     |
| $e^+e^- \rightarrow qq\gamma$                           | 210                              | $1.1 \cdot 10^6$     |
| $e^+e^- \rightarrow qq\gamma\gamma$                     | 47                               | $2.3 \cdot 10^5$     |

Number of signal and background events expected in  $5 ab^{-1}$  of data at 3 TeV CLIC

Since background dominates signal by a factor of  $\sim 700$ , sophisticated event selection is required. It includes:

- Events with 2 isolated photons with  $p_T$  above 15 GeV,  $15 > p_T(\gamma\gamma) > 600$  GeV,  $100 \text{ GeV} > E(\gamma\gamma) > 1000$  GeV
- MVA selection based on 12 sensitive observables in an optimized way to maximize statistical significance.




Reconstructed mass of the selected Higgs boson before (left) and after (right) MVA

- Overall signal selection efficiency  $\sim 43\%$
- Selected number of signal events is 2080 in  $5ab^{-1}$  of integrated luminosity
- Signal to background ratio is 12/1


## 4. Pseudo-experiments

- Experimental data are fully simulated using realistic experimental conditions  $\rightarrow$  "pseudo-data".
- Pseudo-data consisting of signal and background are described by PDF functions ( $f_s$ ,  $f_b$ ) in order to extract the number of signal events  $f = N_s f_s(m_{\gamma\gamma}) + N_b f_b(m_{\gamma\gamma})$ .

- In order to estimate the statistical dissipation of the measured number of signal events, 5000 pseudo-experiments with  $5ab^{-1}$  of data were performed. RMS of the measurement is used as statistical estimator of the uncertainty of the signal count.
- Statistical uncertainty of the Higgs boson to diphoton decay is measured to be 5.5%



One example of pseudo-experiment, showing diphoton invariant mass of pseudo-data (black), corresponding fit with function  $f$  (full line), and background fit with function  $f_b$  (dashed line)



Pull distribution of 5000 pseudo-experiments

## 5. Discussion

- From the fully simulated measurement, we set with the 95% confidence level two tail limit of 10% to the Standard Model prediction of the  $BR(H \rightarrow \gamma\gamma)$ .
- Several sources of systematic uncertainty are considered (uncertainty of photon identification efficiency, uncertainty of integrated luminosity, photon energy resolution, uncertainty of the luminosity spectrum, background modeling). Overall systematic uncertainty is estimated to be  $\sim 2\%$ .
- The obtained results supersedes estimates for 3 TeV CLIC sensitivity obtained from 1.4 TeV simulation luminosity scaling.

This work was published as: G. Kačarević, I. Božović-Jelisavčić, N. Vukašinić, G. Milutinović-Dumbelović, I. Smiljanić, T. Agatonović-Jovin, M. Radulović, and J. Stevanović, Measurement of the Higgs boson branching ratio  $BR(H \rightarrow \gamma\gamma)$  at a 3 TeV CLIC, Phys. Rev. D **105**, 092009, 2022.