

Physics highlights and challenges: particle physics

*Paris Sphicas
CERN & NKUA*

*BPU11 (11th International Conference of the Balkan Physical Union)
Belgrade, Aug 29, 2022*


- ❑ **Introduction: How we got to here: a brief history of the past four decades**
- ❑ **Where we stand & the imminent future (next 10-15 years)**
 - ❑ **The (fundamental?) scalar sector of the SM**
 - ❑ **Direct searches for BSM physics**
- ❑ **Dark Matter**
 - ❑ **Interplay of Direct and Indirect (collider) searches**
- ❑ **Physics of Flavor**
 - ❑ **Quark sector: CP violation, rare decays of K and B mesons**
 - ❑ **Neutrinos: New source of CP violation(?) Mass ordering? Nature? New particles/interactions?**
- ❑ **Pseudo-summary**



H.F.R.I.
Hellenic Foundation for
Research & Innovation

The research project was supported by the Hellenic Foundation for Research and Innovation (H.F.R.I.) under the “1st Call for H.F.R.I. Research Projects to support Faculty Members & Researchers and the Procurement of High-and the procurement of high-cost research equipment grant” (Project Number:2288).

The Standard Model and colliders



Hadron Colliders
W/Z: UA1/UA2 @ SPS
Top: CDF/D0 @ Tevatron
H: ATLAS/CMS @ LHC

Hadron Collisions
b quark: E288 @ FNAL
c quark: pBe @ AGS

ee Colliders
c and τ @ SPEAR (SLAC)
g @ PETRA (DESY)


Not shown: probing/studying the strong, weak and EM interactions

How we got to here

A very (very...) brief summary

pp collisions ::= parton-parton collisions + extra debris


Early 80s: The rendez-vous with the W boson.



It was there, at the right time (number of events → rate → time of rendez-vous!)


At the right mass: 81 ± 5 GeV

With the right spin: 1




The similarly punctual cousin: the Z boson

- **The Z boson was there as well**
 - Also at the right time
 - And at the right mass (91 GeV)



- **Tremendous (other) physics output from the 10 years of the SPS and the UA/UA2 experiments**



End of the 80s, beginning 90s:

Passing the baton to
(a) LEP (CERN) & SLC (Stanford)
and
(b) The Tevatron (Fermilab)


LEP & SLD Overview: Luminosity, Energy, Precision

LEP: Conventional collider e^+e^- ring





Energy upgradeable; measurable;
Four detectors (A,D,L,O); Large luminosity:
20 Million Z events.
But.... Energy limited by synchrotron radiation loss
($\sim \gamma^4$). At Z: 3 GeV/turn (replenished by the RF system)

PEP @ SLAC: complementary to LEP




First (high \sqrt{s}) e^+e^- “linear” collider (+ arcs);
Small transverse beam sizes (& beam pipe)
Reduced luminosity; Only one expt (SLD):
0.35 Million Z events
BUT: 73% electron beam polarization;


e^+e^- collisions: clean, controlled environment



Two-jet event



Three-jet event



The LEP legacy



**LEP ran for over a decade
And delivered unprecedented
precision on the Electroweak Theory
On several observables, 0.1%**

We also learned that:
There are at most three neutrino species
lighter than M_Z . $N_\nu = 2.9919 \pm 0.0081$



**Higgs boson heavier than 114.5 GeV
SUSY not there for masses ≤ 100 GeV**

Highest \sqrt{s} attained: 209.2 GeV.
K. Hubner Phys. Reports 403-404 (2004)
The maximum energy of LEP2 was determined by the decision
in 1996 to discontinue the production of the SC cavities...

Precision gives access to loops





End of 80s, beginning of 90s: Tevatron



The Tevatron discovery: the top quark (I)

- The crowning moment for the Tevatron experiments: the observation of the Top quark
 - The most complicated signature up to that point in time; leptons, jets, missing transverse energy, and b-tagging!




LEP and the Tevatron were huge successes

The word “success” does not do justice
Yet... the Higgs Boson did not show up

And the Higgs... is extremely important:

the entire Standard Model of Particle
Physics is based on the Brout-Englert-
Higgs mechanism to


*give masses to the W & Z bosons, while
leaving the photon massless;
bonus: it also gives masses to the
fermions!*



A machine for EWSB


□ The Large Hadron Collider (LHC)

- Use existing LEP tunnel at CERN
- Replace: e by p ; increase bending power (with superconducting magnets)
- At the time (1990s): mass of the Higgs boson was completely unknown. But theoretical arguments said $M_H < 1 \text{ TeV}$ (1000 GeV).



$$\begin{aligned} M_H &\sim 1000 \text{ GeV} \\ E_W &\geq 500 \text{ GeV} \\ E_q &\geq 1000 \text{ GeV} (1 \text{ TeV}) \\ E_p &\geq 6000 \text{ GeV} (6 \text{ TeV}) \end{aligned}$$

But running at
close to this
energy \rightarrow need
high luminosity
 \rightarrow pileup!




A tremendous, unprecedented challenge




CMS Experiment at the LHC, CERN
Data recorded: 2012-May-13 20:08:14.621490 GMT
Run/Event: 194108 / 564224000

$H \rightarrow \gamma\gamma$







$p_T(\mu) = 36, 48, 26, 72 \text{ GeV}; m_{12} = 86.3 \text{ GeV}$



$H \rightarrow ZZ \rightarrow 4\mu$
 $m_{4\mu} = 125.1 \text{ GeV}$


In July 2012: a new particle with mass $\approx 125 \text{ GeV}$ is seen! It decays to $\gamma\gamma, ZZ$
 \rightarrow it's a boson (spin=0 or 2)






- **Measurement of Spin-Parity, mass: it IS the vacuum particle...**

[illegible]


$J^P=0^+(!!!)$




Phys. Lett. B 784 (2018) 345




(a) Gluon Fusion




(b) Vector Boson Fusion




(c) Higgsstrahlung




(d) $t\bar{t}$ Production



$$\mu = 1.06 \pm 0.07$$





SM @ the highest E; EWSB (“Higgs” sector) (I)




**Clear observation
of couplings to
3rd-gen fermions**

ttH



**Evidence of coupling to
2nd-gen fermions**




CMS:

obs: 3.0 σ (exp: 2.5 σ)

ATLAS:

obs: 2.0 σ (exp: 1.7 σ)

$$\mu = 1.2 \pm 0.6$$



$H \rightarrow c\bar{c}$

$$\mu_{VZ(Z \rightarrow c\bar{c})} = 1.01^{+0.23}_{-0.21}$$

$$\mu_{VH(H \rightarrow c\bar{c})} < 14 \text{ (7.6 exp.)}$$

$$1.1 < |\kappa_c| < 5.5 \text{ (} |\kappa_c| < 3.4 \text{ exp.)}$$


The overall picture of the measured/seen Higgs couplings

$$\lambda_f = \kappa_f \left(\frac{m_f}{v} \right)$$

$$\left(\frac{g_V}{2v} \right)^{1/2} = \kappa_v^{1/2} \left(\frac{m_V}{v} \right)$$

A new kind of “force”,
with *non-universal*
couplings to matter (!)


A particle like no other!



Where we stand now


A very brief summary

Standard Model of Particle Physics



Highest priority:
understand this central
and most strange element

Standard Model Total Production Cross Section Measurements



Putting it all together: SM reigns supreme

Goodness of fit

- $\chi^2_{\min}=18.6 \rightarrow \text{Prob}=23\%$

Fit result often more accurate than measurement


- Small pulls for M_H , M_Z , $\Delta\alpha_{\text{had}}^{(5)}(M_Z^2)$, m_c , $m_b \rightarrow$ input accuracies exceed fit requirements

Knowledge of $m_H \rightarrow$ huge improvement in:

- m_W (28 \rightarrow 11 MeV) arXiv:1407.3792
- m_t (6.2 \rightarrow 2.5 GeV)
- $\sin^2\theta_W$ (2.3 \rightarrow 1.0×10^{-3})

Largest discrepancy:

- $A_{\text{FB}}(b)$: 2.5σ




The discovery of the Higgs boson was the ultimate crown on the Standard Model of Particle Physics

With the discovery of the Higgs boson, the Standard Model (SM) is now “complete”: its full particle content has been observed

The SM provides a remarkably accurate description of experiments with and without high-energy accelerators. At the cost of 26 parameters determined by experiment...

With the physics of the very small [thought to be] understood at energy scales of ≥ 100 GeV, the situation is reminiscent of previous times in history when our knowledge of nature was deemed to be “complete”



Lord Kelvin (1900):

There is nothing new to be discovered in physics now. All that remains is more and more precise measurement.

1905-1920: Relativity, Quantum mechanics...


The magic of the Higgs boson mass

Quantum Mechanics: ultimate destructor of small numbers (in nature) not protected by some symmetry (thus “law”)

Higgs boson: the ultimate example; spinless \rightarrow zero cost from mass correction

$$m^2(p^2) = m_o^2 + \text{[wavy line with } J=1 \text{]} + \text{[circle with } J=1/2 \text{]} + \text{[loop with } J=0 \text{]}$$

$$m^2(p^2) = m^2(\Lambda^2) + Cg^2 \int_{p^2}^{\Lambda^2} dk^2$$



P.A.M Dirac


M(H) with corrections all the way up to the Planck scale: for $\Lambda \sim 10^{19}$ GeV

[for illustration only; numbers are random]

$$m^2 = 1234567890123456789012345675432189012 - 1234567890123456789012345675432173387 = 15625 \text{ GeV}^2$$

An immense coincidence of googlic sizes?
Probably, simply some additional (i.e. New) physics on the way to 10^{19} GeV

We know there is new physics already



Dark (invisible) matter!

Plus neutrinos and their masses!

The hunt for “New Physics”

**Despite finding the last missing
piece of the SM, there is plenty of
room for NP**

(beyond the magic of the M_H - m_t mass)

**Some real and some virtual reasons to believe in
new physics**

Real reasons: dark matter & ν masses

Virtual reasons: naturalness

+ matter–antimatter asymmetry in the universe...


Solutions to the hierarchy problem

□ Solution #1: a composite Higgs

- H: bound state (e.g. due to some new strong interaction)

□ Solution #2: supersymmetry

- Partners for ALL SM particles, spin different by $\frac{1}{2}$
- Presumably broken symmetry (since partners unseen)



□ Solution #3: little Higgs

- H: pseudo-Goldstone boson of Ultimate Theory; just another effective theory, e.g. valid to ~ 10 TeV. Loops cancel by particles of same spin (so need to introduce these particles)


□ Solution #4: extra dimensions

- N-dim space; gravity propagates in all dims, SM only in “our” 3 dims; e.g. warped extra dimension can explain weakness of G
- All of these “solutions” introduce either deviations in tails or new particles, with masses whose “natural” values should be O(few TeV)


What we have been looking for (mainly)

CMS Experiment at LHC, CERN
Data recorded: Wed Jun 13 21:51:54 2012 PDT
Run/Event: 196250 / 615309469
Lumi section: 385
Orbit/Crossing: 100914566 / 2074

HT = 1009 GeV



CMS Experiment at LHC, CERN
Data recorded: Tue Jun 14 11:40:21 2011 PDT
Run/Event: 166966 / 147335241
Lumi section: 169




The never-ending search for SUSY... (and it's not like we didn't look for them)

ATLAS SUSY Searches* - 95% CL Lower Limits

June 2021

ATLAS Preliminary

$\sqrt{s} = 13$ TeV




*Only a selection of the available mass limits on new states or phenomena is shown. Many of the limits are based on simplified models, c.f. refs. for the assumptions made.

10⁻¹ 1 Mass scale [TeV]

The High-Luminosity LHC (HL-LHC)



HL-LHC in a nutshell
Some physics projections

From the LHC to High-Luminosity LHC (HL-LHC)



Today

HL-LHC: major intervention on more than 1.2 km of the LHC with new technologies: Nb₃Sn magnets, Crab cavities,...




Last updated: January 2022

HL-LHC challenges

Annual dose at HL-LHC:
similar to total dose from
LHC start to LS3


Key to physics: maintain
detector performance in
the presence of much
higher pileup (140-200!)



Upgrade several detector components (trackers, calorimeters, redesign some electronics, new detector technologies, Trigger and DAQ)


Medium-term Higgs physics: the LHC/HL-LHC program

HL-LHC reach: Higgs






H width to invisible:
 $h(125) \rightarrow XX$.

Includes BSM decays
and rare SM decays: $\leq 4\%$



The ultimate frontier: Higgs self-coupling

$$V_h = \frac{m_h^2}{2} h^2 + (1 + \kappa_3) \lambda_{hhh}^{\text{SM}} v h^3 + \frac{1}{4} (1 + \kappa_4) \lambda_{hhhh}^{\text{SM}} h^4$$




Long-term future: proposed machines

A very brief summary

EU Strategy: High-priority future initiatives

- ❑ **A. An electron-positron Higgs factory is the highest-priority next collider. For the longer term, the European particle physics community has the ambition to operate a proton-proton collider at the highest achievable energy. Accomplishing these compelling goals will require innovation and cutting-edge technology:**
 - ❑ *the particle physics community should ramp up its R&D effort focused on advanced accelerator technologies, in particular that for high-field superconducting magnets, including high-temperature superconductors;*
 - ❑ *Europe, together with its international partners, should investigate the technical and financial feasibility of a future hadron collider at CERN with a centre-of-mass energy of at least 100 TeV and with an electron-positron Higgs and electroweak factory as a possible first stage. Such a feasibility study of the colliders and related infrastructure should be established as a global endeavour and be completed on the timescale of the next Strategy update.*
 - ❑ *The timely realisation of the electron-positron International Linear Collider (ILC) in Japan would be compatible with this strategy and, in that case, the European particle physics community would wish to collaborate.*

Collider Crib sheet



□ ILC (Japan):


- Linear collider with high-gradient superconducting acceleration
- Ultimate: 0.5-1(?) TeV
- To secure (...) funding: reduce cost by starting at 250 GeV (H factory)

□ CLIC (CERN):

- Linear collider with high gradient normal-conducting acceleration
- Ultimate: multi-TeV (3) e^+e^- collisions
- Use technology to overcome challenges
- Stages, for physics and funding

□ FCC-ee/FCC-hh (CERN):

- Protons to extend energy frontier
- 100 km ring with 16T magnets
- Use FCC-hh tunnel for e^+e^- collider
- Technology for ee: “standard”









□ CEPC/SppC

- Essentially an FCC-ee, then hh with (a) more conservative luminosity estimates and (b) in China

□ Outliers:


- “Low-field” (7T) magnets @ FCC (?)
- Muon Collider (???)

The Higgs sector: from the HL-LHC to the “future”




Precision Observables & Searches: examples


- EWPO:
circular ee
colliders
+ linear colliders
for $\sin^2\theta_W$.
Note: currently,
discussion/plan
for a large Z run
for the linear
colliders...



Higgs compositeness (?)




Electroweak SUSY reach





EWPO	Current	CEPC	FCC (ee)
M_Z [MeV]	2.1	0.5	0.1
Γ_Z [MeV]	2.1	0.5	0.1
N_ν [%]	1.7	0.05	0.03
M_W [MeV]	12	1	0.67
$A_{FB}^{0,b}$ [$\times 10^4$]	16	1	< 1
$\sin^2\theta_W^{\text{eff}}$ [$\times 10^5$]	16	1	0.6
R_b^0 [$\times 10^5$]	66	4	2–6
R_μ^0 [$\times 10^5$]	2500	200	100

The Dark Sector

- ❑ **An experimental fact & yet, still a total mystery**
 - ❑ And masses span over 80 orders of magnitude
- ❑ **Nightmare scenario: totally dark**
 - ❑ Only Gravity to play with...




- ❑ **More promising: some shade of grey**




DM: direct detection experiments

Two-phase liquid-Xe
expts continuing to
increase their
sensitivity





L. Hsu ICHEP2020



Very large program for subGeV DM

L. Hsu ICHEP2020




HeRALD




DM: Classic WIMPs

- Two (SUSY) “extremes”, pure Wino, pure Higgsino
 - Main “tools”: disappearing track, propagator modifications



For small Δm , soft π^\pm ...




EWKinOs in loop change prop
(W, Y parameters)



Scalar mediator: Higgs portal and BSM scalar



A collider discovery will need confirmation from DD/ID for cosmological origin
 A DD/ID discovery will need confirmation from colliders to understand the nature of the interaction



A future collider program that optimizes sensitivity to invisible particles coherently with DD/ID serves us well. Need maximum overlap with DD/ID!

Physics of Flavor

Neutrinos

Quarks

Charged Leptons

Super-brief Intro: from hopeless to lucky strike(s)

From invisible particle ($\sigma \sim 10^{-44} \text{cm}^2$ @ $E_\nu \sim 2 \text{ MeV}$) to major source of physics:

From Pontecorvo's few/day/ton with $\sim 10^{11} \nu/\text{cm}^2/\text{s}$ to \rightarrow PMNS matrix & CP violation(?!?)

$$P(\nu_\alpha \rightarrow \nu_\beta) = \sin^2 2\theta \sin^2 \left(1.27 \frac{\Delta m^2 (\text{eV}^2) L (\text{km})}{E_\nu (\text{GeV})} \right) \quad P(\nu_\alpha \rightarrow \nu_\alpha) = 1 - P(\nu_\alpha \rightarrow \nu_\beta)$$

SuperK, SNO: large mixing; good Δm
 $\Delta m_{\text{sol}}^2 \approx 7 - 8 \times 10^{-5} \text{eV}^2$

$\Delta m_{\text{sol}}^2 \sim E_\nu (\text{MeV}) / L (100 \text{ km})$
 \rightarrow Reactors! \rightarrow KamLAND

IMB, Kamiokande, Soudan(2)... SuperK:
 large mixing; super Δm : $\Delta m_{\text{atm}}^2 \approx 2.5 \times 10^{-3} \text{eV}^2$

$\Delta m_{\text{atm}}^2 \sim E_\nu (\text{MeV}) / L (100 \text{ m}) = (\text{GeV}) / (100 \text{ km})$
 \rightarrow Accelerators! \rightarrow K2K, MINOS(+), ..., T2K, NOVA
 + Short baselines: Daya Bay, RENO, ..., xBOONE


ν physics: PMNS; 3x3 or 4x4? nature (Majorana or Dirac); mass ordering

$$U_{3 \times 3} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \cos \theta_{23} & \sin \theta_{23} \\ 0 & -\sin \theta_{23} & \cos \theta_{23} \end{pmatrix} \begin{pmatrix} \cos \theta_{13} & 0 & \sin \theta_{13} e^{-i\delta} \\ 0 & 1 & 0 \\ -\sin \theta_{13} e^{i\delta} & 0 & \cos \theta_{13} \end{pmatrix} \begin{pmatrix} \cos \theta_{12} & \sin \theta_{12} & 0 \\ -\sin \theta_{12} & \cos \theta_{12} & 0 \\ 0 & 0 & 1 \end{pmatrix} \times (U_{\text{Maj}})$$

atmospheric +
 accelerator disapp


SBL reactor +
 accelerator app

solar +
 KamLAND




Neutrinos: Reactor experiments; Precision!

Principle: θ_{13}




The prized measurement



Third lucky strike (in ν phys)

$$J_{CP}^{\text{lep}} = \text{Im}(U_{\mu 3} U_{e 3}^* U_{e 2} U_{\mu 2}^*) = (1/8) \times \\ \times \sin 2\theta_{12} \sin 2\theta_{23} \sin 2\theta_{13} \cos \theta_{13} \sin \delta_{CP} \\ \approx 0.033 \sin \delta_{CP} \text{ BUT : } J_{CP}^q \approx 3 \times 10^{-5}$$

Long series of measurements have brought us to a “precision era”



de Salas et al, arXiv:2006.11237

M. Tortola ICHEP2020

parameter	best fit $\pm 1\sigma$	3σ range
Δm_{21}^2 [10^{-5}eV^2]	$7.50^{+0.22}_{-0.20}$	6.94–8.14
$ \Delta m_{31}^2 $ [10^{-3}eV^2] (NO)	$2.56^{+0.03}_{-0.04}$	2.46–2.65
$ \Delta m_{31}^2 $ [10^{-3}eV^2] (IO)	2.46 ± 0.03	2.37–2.55
$\sin^2 \theta_{12} / 10^{-1}$	3.18 ± 0.16	2.71–3.70
$\sin^2 \theta_{23} / 10^{-1}$ (NO)	$5.66^{+0.16}_{-0.22}$	4.41–6.09
$\sin^2 \theta_{23} / 10^{-1}$ (IO)	$5.66^{+0.18}_{-0.23}$	4.46–6.09
$\sin^2 \theta_{13} / 10^{-2}$ (NO)	$2.225^{+0.055}_{-0.078}$	2.015–2.417
$\sin^2 \theta_{13} / 10^{-2}$ (IO)	$2.250^{+0.056}_{-0.076}$	2.039–2.441
δ/π (NO)	$1.20^{+0.23}_{-0.14}$	0.80–2.00
δ/π (IO)	1.54 ± 0.13	1.14–1.90

2.7%

1.2%

5.2%

4.9%

4.8%

3.0%

17%

8%

relative 1σ uncertainty

3-5%


Neutrinos: oscillations & CPV

Currently (NOvA and T2K) a bit confused


$$\begin{aligned}
 P(\nu_\mu \rightarrow \nu_e) = & \sin^2 \theta_{23} \frac{\sin^2 2\theta_{13}}{(A-1)^2} \sin^2[(A-1)\Delta_{31}] \\
 & + \alpha^2 \cos^2 \theta_{23} \frac{\sin^2 2\theta_{12}}{A^2} \sin^2(A\Delta_{31}) \\
 & - \alpha \frac{\sin 2\theta_{12} \sin 2\theta_{13} \sin 2\theta_{23} \cos \theta_{13} \sin \delta_{CP}}{A(1-A)} \sin \Delta_{31} \sin(A\Delta_{31}) \sin[(1-A)\Delta_{31}] \\
 & + \alpha \frac{\sin 2\theta_{12} \sin 2\theta_{13} \sin 2\theta_{23} \cos \theta_{13} \cos \delta_{CP}}{A(1-A)} \cos \Delta_{31} \sin(A\Delta_{31}) \sin[(1-A)\Delta_{31}]
 \end{aligned}$$

Favored (slightly):
Normal Ordering &
 $\delta_{CP} \sim 3\pi/2$

But... fits are complicated




Unknown
mass
order






T2K-II

NormalO

Known
mass
order



T2K-II



Very useful/nice: complementarity of NOvA/T2K and DUNE/HyperK:
baseline (1300 vs 200 km) and detection technique


Flavor physics: quark sector

$$V_{\text{CKM}} = \begin{bmatrix} 1 - \lambda^2/2 & \lambda & A\lambda^3(\rho - i\eta) \\ -\lambda & 1 - \lambda^2/2 & A\lambda^2 \\ A\lambda^3(1 - \rho - i\eta) & -A\lambda^2 & 1 \end{bmatrix}$$


$$V_{ui} V_{uj}^* + V_{ci} V_{cj}^* + V_{ti} V_{tj}^* = 0$$

CKM triangle: extensive precision program en route



Today





2025



2035



CP $\sim \sin 2\phi$



$$\frac{\Gamma(\bar{B}^0 \rightarrow \bar{f}) - \Gamma(B^0 \rightarrow f)}{\Gamma(\bar{B}^0 \rightarrow \bar{f}) + \Gamma(B^0 \rightarrow f)} = -\eta_f \sin(2\phi) \sin(\Delta M t)$$

Pseudo-summary


Pseudosummary/Outlook

- ❑ **Extremely rich physics program ahead to understand the scalar sector**
 - ❑ The LHC and HL-LHC will get us to 3-5% couplings for the Higgs boson;
 - ❑ All options for a future Higgs “factory” bring in $\sim O(10^{-2}-10^{-3})$ understanding of couplings.
 - ❑ Important aspects: EWPO (needs next-gen Z factory) and top threshold.
 - ❑ **Actually, the linear and circular options are quite complementary...**
 - ❑ Fundamental scalar? Can probe it to 15-18 TeV.
 - ❑ FCC-ee/hh combination has the largest direct reach to new particles/phenomena. From new particles to Higgs self-coupling to Dark Matter searches...
- ❑ **Dark Matter:**
 - ❑ Complementarity with indirect searches at colliders (and astroparticle expts); Next-gen colliders can cover the thermal WIMP scenario.
- ❑ **Flavor Physics: neutrinos have all one could ask for (albeit it with small σ_{int}). But quark sector may hold the first genuine surprises(?)**
 - ❑ In the next 5-10 years: could get very pleasant news on CP front. Definitive statement of mass hierarchy and CP from the full program (DUNE and HyperK).
 - ❑ Quark sector: the current situation will be resolved in the next five years.
- ❑ **Physics at hand, physics of the next decade, physics of the long-term future: remains fully exciting. Stay tuned!**

Backups

Higgs boson issues


- ❑ The only (?!) spin-0 fundamental (?) particle
- ❑ What creates the infamous couplings to the leptons?
 - ❑ It's a new interaction! The only non-universal one (it's not a gauge interaction...), with a *free* parameter (Yukawa coupling) for each combination (worse for quarks – mass matrix...)
 - ❑ What about them neutrinos?!?
- ❑ What protects its mass and sets it to the EWK scale when it should be at Λ_{pl} ?
 - ❑ There is only one dimensionfull variable in the SM Lagrangian: $v=246$ GeV (...!?!...)
 - ❑ The rest of the SM is scale-free...
- ❑ And.... where is all that vaccum energy?
Cosmological constant is $> 10^{100}$ times off.



“Turn on the LHC and... find Higgs & SUSY”

- ❑ **ATLAS and CMS were designed to do this; they were “guaranteed” to find the Higgs – period; right away**
 - ❑ **In fact: SUSY is strongly produced, so will be observed first**
 - ❑ **For the “impatient”: join SUSY physics group**


- ❑ **Many hard Jets**
- ❑ **Large missing energy**
 - ❑ **2 LSPs**
 - ❑ **Many neutrinos**
- ❑ **Many leptons**
- ❑ **In a word Spectacular!**



Text & simu from 1999

The magic of the Higgs boson mass

- **Quantum Mechanics: ultimate destructor of small numbers (in nature) that are not protected by some symmetry (thus “law”)**
- **Higgs boson: the ultimate example**
 - **Quadratic divergence in the Higgs mass**




$$m^2(p^2) = m^2(\Lambda^2) + Cg^2 \int_{p^2}^{\Lambda^2} dk^2$$

- **H mass should be $\sim 10^{19}\text{GeV}$**
 - **Yet, it lies at 125 GeV...**
- **Put differently: if cut off at Λ_{PL} , why $m_W \ll M_{\text{Pl}}$?**
 - **Or, why is gravity ($G \sim 1/M_{\text{Pl}}$) so weak?**

- **Reminder of just two past applications of naturalness argument:**
- **Weisskopf (1939): “the self-energy of charged particles obeying Bose statistics is found to be quadratically divergent...,”**
 - in theories of elementary bosons, new phenomena must enter at an energy scale of m/e
 - + positron, doubling of particles...

- **Rare Kaon decays: $K_L \rightarrow \mu^+ \mu^-$**



GIM proposal

$$\sim G \Lambda^2 = G(m_c^2 - m_u^2)$$

$$\rightarrow \Lambda^2 \sim 3-4 \text{ GeV}^2 !!!$$

Other open issues in particle physics

- ❑ **Is the Higgs boson at 125 GeV the Higgs boson of the Standard Model? Are there other Higgs-like bosons?**
- ❑ **How did (does) the current matter–antimatter asymmetry in the universe arise?**
- ❑ **There is now experimental evidence (astrophysics!) that there is a mysterious type of matter: “Dark” matter. What is it? What does it interact with?**
- ❑ **Neutrinos (which are taken as massless) turned out to have masses. But tiny-tiny masses. Natural explanation of this smallness is New Physics**
- ❑ **And while we’re at it: gravity remains totally intractable (beyond astrophysics that is...)**






The road to obtaining answers: precise measurements & direct searches...

- ❑ **Physics of fundamental scalars and direct searches for New Physics**
 - ❑ LHC, HL-LHC & future collider(s) (FCC-ee, FCC-hh, FCC-eh, CEPC, ILC, CLIC)
 - ❑ Understand the strangest of all elements of the SM.
 - ❑ Search for new particles and physics phenomena.
- ❑ **Dark Matter searches**
 - ❑ Very large number of experiments aiming at detecting DM
 - ❑ Direct detection (lab), indirect detection (astrophy).
 - ❑ Matter-matter collisions.
- ❑ **Physics of Flavor:**
 - ❑ Quark sector: Kaon and B hadron decays (LHC, HL-LHC, Fixed-Target (CERN, JP), Belle II (JP))
 - ❑ Measure CP sources as precisely as possible. Probe sacred (but also accidental...) laws (e.g. lepton number conservation). Universality? Probe rarest decay modes.
 - ❑ Neutrinos: Complementarity of beams [DUNE (US), HyperK (JP)], reactors (JUNO), atmospheric (ORCA)
 - ❑ Is there CP violation in the lepton sector? PMNS matrix (à la CKM)?
 - ❑ Mass hierarchy of three ν generations (“normal” or “inverted”)? Dirac or Majorana?
 - ❑ High-precision measurements; tests of Fundamental Symmetries

Future accelerators

Collider	Type	\sqrt{s}	\mathcal{P} [%] [e^-/e^+]	N(Det.)	$\mathcal{L}_{\text{inst}}$ [10^{34}] $\text{cm}^{-2}\text{s}^{-1}$	\mathcal{L} [ab^{-1}]	Time [years]	Refs.	Abbreviation
HL-LHC	pp	14 TeV	-	2	5	6.0	12	[10]	HL-LHC
HE-LHC	pp	27 TeV	-	2	16	15.0	20	[10]	HE-LHC
FCC-hh	pp	100 TeV	-	2	30	30.0	25	[1]	FCC-hh
FCC-ee	ee	M_Z	0/0	2	100/200	150	4	[1]	FCC-ee ₂₄₀ FCC-ee ₃₆₅ (1y SD before $2m_{\text{top}}$ run)
		$2M_W$	0/0	2	25	10	1-2		
		240 GeV	0/0	2	7	5	3		
		$2m_{\text{top}}$	0/0	2	0.8/1.4	1.5	5 (+1)		
ILC	ee	250 GeV	$\pm 80/\pm 30$	1	1.35/2.7	2.0	11.5	[3, 11]	ILC ₂₅₀ ILC ₃₅₀ ILC ₅₀₀ (1y SD after 250 GeV run)
		350 GeV	$\pm 80/\pm 30$	1	1.6	0.2	1		
		500 GeV	$\pm 80/\pm 30$	1	1.8/3.6	4.0	8.5 (+1)		
CEPC	ee	M_Z	0/0	2	17/32	16	2	[2]	CEPC
		$2M_W$	0/0	2	10	2.6	1		
		240 GeV	0/0	2	3	5.6	7		
CLIC	ee	380 GeV	$\pm 80/0$	1	1.5	1.0	8	[12]	CLIC ₃₈₀ CLIC ₁₅₀₀ CLIC ₃₀₀₀ (2y SDs between energy stages)
		1.5 TeV	$\pm 80/0$	1	3.7	2.5	7		
		3.0 TeV	$\pm 80/0$	1	6.0	5.0	8 (+4)		
LHeC	ep	1.3 TeV	-	1	0.8	1.0	15	[9]	LHeC
HE-LHeC	ep	1.8 TeV	-	1	1.5	2.0	20	[1]	HE-LHeC
FCC-eh	ep	3.5 TeV	-	1	1.5	2.0	25	[1]	FCC-eh

ILC, CLIC, FCC-ee/hh, CEPC



CEPC: multiple candidate sites in China

The challenges

□ Linear ee colliders:

- Acceleration gradient
 - ILC: 30 MV/m; CLIC: 72 MV/m
- Luminosity (to be partially recovered by polarization)
 - + brem loss: e.g. at 3000 GeV, 1/3 at $>0.99\sqrt{s}$
- Tiny beam spot:
 - v : 8 nm for ILC; 3 nm for CLIC3000.
- Power consumption
 - ILC: 130-300 MW
 - CLIC: 170-590 MW

□ Circular ee colliders:

- Power consumption: 260-350 MW
- Luminosity drops with E

□ hh collider:

- Magnets!
 - Need 16 TeV (x2 LHC); they do not exist today
- High stored beam energy (8-9 GJ)
 - Beam handling, beam dumping
 - Collimation
- High synchrotron radiation inside magnets: several MW
- Beam screen design and cryogenic efficiency;
- Power consumption: 580 MW

Costs (GUnits):

ILC: 5.0 (for 250 GeV); 7.8 (500 GeV)





CLIC: 5.9 or 7.3 (for 380 GeV) + 5.1 (1500 GeV) + 7.1 (3000 GeV) (Tot: 19.5)

FCC-ee: 11.6; but 7.1 is the tunnel

FCC-hh: tunnel + 17 (Tot: 24)

FCC-ee and CEPC

- ❑ **Double-ring colliders with full-energy top-up booster ring**
 - ❑ CEPC started as 54 km, single-ring design; nowadays ~ FCC-ee 100 km, double-ring
 - ❑ 2 IPs, 2 RF straights, tapering of arc magnet strengths to match local energy
 - ❑ Common use of RF systems for both beams at highest energy working point
 - ❑ Synchrotron radiation: 50 MW (30 MW) at FCC-ee (CEPC)
 - ❑ Beam lifetime >12 min; top-up injection, e^+ rate $\sim 10^{11}/s$.
- ❑ **Asymmetric IRs: limit SR of incoming beams towards detectors**
 - ❑ → large crossing angle




FCC-hh and SppC

- **Circumference ~100 km, high luminosity $3(1) \times 10^{35} \text{cm}^{-2}\text{s}^{-1}$ (SppC)**
 - Two IRs at high lumi potentially two more experiments (possibly combined with injection section, collimation insertions, extraction/dump insertion, RF insertion,


FCC-hh

based on
existing CERN
injector chain,

Luminosity goal
~20 ab^{-1} per
main IP within
25 years



SppC



new injector
chain,
simultaneous
operation with
 e^+e^- collider

Grand summary


kappa-3 scenario	ILC ₂₅₀	ILC ₅₀₀	CLIC ₃₈₀	CLIC ₁₅₀₀	HL-LHC+ CLIC ₃₀₀₀	CEPC	FCC-ee ₂₄₀	FCC-ee ₃₆₅	FCC-ee/eh/hh
κ_W (%)	1.1	0.29	0.75	0.4	0.38	0.95	0.95	0.41	0.2
κ_Z (%)	0.29	0.23	0.44	0.39	0.39	0.18	0.19	0.17	0.17
κ_g (%)	1.4	0.84	1.5	1.1	0.86	1.1	1.2	0.89	0.53
κ_γ (%)	1.3	1.2	1.5*	1.3	1.1	1.2	1.3	1.2	0.36
$\kappa_{Z\gamma}$ (%)	11.*	11.*	11.*	8.4	5.7	6.3	11.*	10.	0.7
κ_c (%)	2.	1.2	4.1	1.9	1.4	2.	1.6	1.3	0.97
κ_t (%)	2.7	2.4	2.7	1.9	1.9	2.6	2.6	2.6	0.95
κ_b (%)	1.2	0.57	1.2	0.61	0.53	0.92	1.	0.64	0.48
κ_μ (%)	4.2	3.9	4.4*	4.1	3.5	3.9	4.	3.9	0.44
κ_τ (%)	1.1	0.64	1.4	0.99	0.82	0.96	0.98	0.66	0.49
BR _{inv} (<%, 95% CL)	0.26	0.22	0.63	0.62	0.61	0.27	0.22	0.19	0.024
BR _{unt} (<%, 95% CL)	1.8	1.4	2.7	2.4	2.4	1.1	1.2	1.	1.

All ee colliders achieve major (and comparable) improvements in their first stage already in probing Higgs sector compared to HL-LHC: at least half of the couplings get improved by factor 5 or more W/Z effective couplings and BR(H → invisible) probed to $\sim 3 \times 10^{-3}$


Model-independent total cross section measurement → access to width, untagged BR

Clean environment to study H if/when anomalies are seen to understand underlying physics


Grand summary: Higgs couplings (κ framework) (Future Collider+HL-LHC)



Plus of κ framework: it is simple; Minus: underestimates effects of polarization
Can show deviation from SM, but no real further information on nature of source of deviation;
Untagged and invisible BRs constrained by measurements.




Higgs “factories”



Schemes for increasing luminosity:


- FCC-ee: consider more IRs/running longer
- ILC: more bunches per pulse, doubling repetition rate?
 - Each: x 2 in lumi; higher power consumption and somewhat higher cost
- CLIC: doubling repetition rate at 380 GeV?
 - Factor 2 in lumi; power increases from 170 MW to 220 MW (+slight cost increase)



Low energies: circular colliders superior
 performance Higher energies: CC lumi reduction
 due to synchrotron radiation; linear colliders better:
 luminosity per beam power roughly constant

Longit. polarisation: only at Linear Colliders
 e^- : 80%; e^+ : 30% @ ILC; 0 @ CLIC (not needed)
 FCC-ee: transverse polarization for precise E_{beam}

Higgs Compositeness?



$$\text{gauge } \rho \sim \frac{g_W}{g_*}$$

H


$$\frac{c_\phi}{\Lambda^2} = \frac{g_*^2}{m_*^2}$$

2f-2b

$$\frac{c_W}{\Lambda^2} = \frac{1}{m_*^2}$$

4f, W'/Z'


$$\frac{c_{2W}}{\Lambda^2} = \frac{1}{g_*^2 m_*^2}$$



O_ϕ (Higgs measurements, primarily FCC-ee & CLIC)

Direct resonance searches (generic “ ρ ” EWK triplet; dileptons & dibosons)


Obtain limit on m_* , i.e. on $1/\text{length}(H)$



New resonances/particles/forces?

Seeing the peak. Reach:


- $M < \sqrt{s}$ for lepton colliders
- $M \lesssim 0.3-0.5 \sqrt{s}$ in hadron colliders for couplings \sim weak couplings





Courtesy:
J. De Blas

Deviations in high-M tails:

- Better suited for lepton colliders; sensitive to $[mass/coupling] \gg \sqrt{s}$
- Hadron colliders relevant for $g_{Z'} > g_{SM}$ couplings: $[mass/coupling] \gg 0.5\sqrt{s}$




Use very simple model as example. Universal Z' . Clearly, many models with flavor dependence etc.



Higgs Compositeness?

Using fits from EWK/Higgs group ([arXiv:1905.03764](https://arxiv.org/abs/1905.03764))

Connection between notations:



$$\frac{c_\phi}{\Lambda^2} = \frac{g_*^2}{m_*^2}$$

H

$$\frac{c_W}{\Lambda^2} = \frac{1}{m_*^2}$$

2f-2b


$$\frac{c_{2W}}{\Lambda^2} = \frac{1}{g_*^2 m_*^2}$$

4f, W'/Z'

$$\frac{\text{gauge } \rho}{\Lambda^2} \sim \frac{g_W}{g_*}$$

For mass/coupling ~ 2 TeV \rightarrow deviations $\sim 1\%$ in Higgs couplings

95% CL limits on compositeness scale (O_H operator)



**Maximum sensitivities:
@CLIC and
FCC(ee+eh+hh)**

Long-term future: SUSY at >1TeV.... <10 TeV?


□ The questions:

- If {SUSY} which masses (and mass differences) of strongly- or weakly-coupled superpartners can we reach?
- Is nature fundamentally fine-tuned? If the solution is SUSY, how well can we test this?
- Is dark matter a thermal SUSY WIMP?

□ Strongly-interacting SUSY (gluinos and squarks): simply, the purview of hadron colliders


Hadron Colliders: gluino projections

(R-parity conserving SUSY, prompt searches)



HE-LHC: $\sim 2 \times M_{\text{HL-LHC}}$
FCC-hh: $\sim 5 \times M_{\text{HL-LHC}}$

All Colliders: Top squark projections (R-parity conserving SUSY, prompt searches)



ILC 500: discovery in all scenarios up to kinematic limit $\sqrt{s}/2$


Long-term future: SUSY EWK sector

ee: mainly $\chi_2^0 \chi_1^0$ and $\chi_1^+ \chi_1^-$ ($\Delta m > 1 \text{ GeV}$)
 pp: mainly $\chi_2^0 \chi_1^\pm$ (gaps at low Δm)



ee sensitivity
 ~independent
 of nature of
 LSP
 (but up to $\sqrt{s}/2$)

Thermal
 WIMP

pp sensitivity:
 depends on LSP;
 ISR jet for very low
 Δm (up to $\sim 5 \text{ GeV}$);
 Pure Higgsino:
 stopped track





Higgsino-like EWK processes



Scalar mediator: Higgs portal and BSM scalar

A collider discovery will need confirmation from DD/ID for cosmological origin

A DD/ID discovery will need confirmation from colliders to understand the nature of the interaction




A future collider program that optimizes sensitivity to invisible particles coherently with DD/ID serves us well. Need maximum overlap with DD/ID!


Extended Scalar sectors: MSSM

Simply: hadron colliders...


LHC+HL-LHC



$pp \rightarrow bbH^0/A \rightarrow bb\tau\tau$ (large $\tan\beta$)
 $pp \rightarrow bbH^0/A \rightarrow ttbb$ (int. $\tan\beta$)
 $pp \rightarrow ttH^0/A \rightarrow tttt$ (low $\tan\beta$)



$pp \rightarrow btH^\pm \rightarrow bb\tau\nu$
 $pp \rightarrow btH^\pm \rightarrow tbtb$



Indirect info also probes additional h bosons (e.g. $\kappa_b \sim m_Z^2/m_A^2$)

H^0/A : exclusion limits > 5 TeV
 (20 TeV at low $\tan \beta$)
 H^\pm : exclusion limits $\sim 10 - 15$ TeV

EU Strategy: other essential scientific activities for particle physics

- ❑ **A. The quest for dark matter and the exploration of flavour and fundamental symmetries are crucial components of the search for new physics.**
 - ❑ This search can be done in many ways, for example through precision measurements of flavour physics and electric or magnetic dipole moments, and searches for axions, dark sector candidates and feebly interacting particles. There are many options to address such physics topics including energy-frontier colliders, accelerator and non-accelerator experiments. A diverse programme that is complementary to the energy frontier is an essential part of the European particle physics Strategy. *Experiments in such diverse areas that offer potential high-impact particle physics programmes at laboratories in Europe should be supported, as well as participation in such experiments in other regions of the world.*
- ❑ **B. Theoretical physics is an essential driver of particle physics that opens new, daring lines of research, motivates experimental searches and provides the tools needed to fully exploit experimental results. It also plays an important role in capturing the imagination of the public and inspiring young researchers.**
- ❑ **C. The success of particle physics experiments relies on innovative instrumentation and state-of-the-art infrastructures. To prepare and realise future experimental research programmes, the community must maintain a strong focus on instrumentation.**

Where we stand


- ❑ **Most successful Theory ever: Standard Model of Particle Physics**
 - ❑ **Highest priority:** *extend understanding of SM and its newly discovered scalar sector.*
- ❑ **Evidence that SM is either incomplete or an Effective Theory of some Ultimate Theory (or another step in a series of Effective Theories...)**
 - ❑ **Experimental evidence**
 - ❑ **Dark Matter (DM):** The “thing” we know the least about: Unknown nature, unknown number(s) of DM particles, unknown mass range(s) – 10^{80} .
 - ❑ **Neutrino masses:** SM gauge group allows for Majorana masses, “explaining” their tiny values; unknown (putative) Majorana scale.
 - ❑ **Matter-antimatter asymmetry of the Universe:** nowhere near what we measure in CP violation experiments. What about CP violation in the lepton sector?
 - ❑ **Theory issues**
 - ❑ **Electroweak (EW) hierarchy problem.** Why is the Higgs so light?
 - ❑ **The flavour puzzle.** Why three generations of quarks and leptons? With very different masses and mixings? Size of CP violation? (explain matter universe???)
 - ❑ **And lots more...** e.g. Strong CP problem. Another vacuum issue (this time QCD). Why is its θ parameter experimentally constrained to be extremely small? For a priori no good reason.... → Axions?

SM Outliers


Current issues to watch...

A very sensitive SM test: the m_W – m_t plane



W mass



$$m_W^2 \left(1 - \frac{m_W^2}{m_Z^2} \right) = \frac{\pi\alpha}{\sqrt{2}G_\mu} (1 + \Delta r)$$




t mass



A very sensitive SM test: the m_W – m_t plane



W mass



$m_W(\text{SM}) = 80375 \pm 4(\text{inp}) \pm 4(\text{th}) \text{ MeV}$
 $m_W(\text{CDF}) = 80433 \pm 9.4 \text{ MeV}$
 $\Delta m_W(\text{SM-CDF}) = 7\sigma \dots$


More precise than all previous m_W measurements combined
 ~ 3σ higher than most precise other measurements
 ~ 3σ higher than previous CDF result (1/4 of data, correlations taken into account)

$$m_W^2 \left(1 - \frac{m_W^2}{m_Z^2} \right) = \frac{\pi\alpha}{\sqrt{2}G_\mu} (1 + \Delta r)$$



“in tension” with SM pred

Muon $a_\mu = g_\mu - 2$





$$a_\mu^{\text{QED}} = \frac{\alpha}{2\pi} + A_2 \left(\frac{\alpha}{\pi}\right)^2 + A_3 \left(\frac{\alpha}{\pi}\right)^3 + A_4 \left(\frac{\alpha}{\pi}\right)^4 + A_5 \left(\frac{\alpha}{\pi}\right)^5$$

$$a_\mu^{\text{EW}} = 153.6 (1.0) \times 10^{-11}$$





$$a_\mu^{\text{HLbL}} = 94 (19) \times 10^{-11}$$

Main uncertainty: Hadronic Vacuum Polarization




Low-energy e^+e^- or Lattice QCD

4.2 σ ; but:





Flavor physics: charged lepton sector & summary

Final MEG upper limit
 $B(\mu \rightarrow e\gamma) < 4.2 \cdot 10^{-13}$
 @90% CL: there
 since 2016...




Muon LFV quiet since 2016...
 Exciting times ahead with
 MEGII, Mu2e, COMET, Mu3e...

$\mu \rightarrow e\gamma$
 $\mu \rightarrow eee$
 $\mu N \rightarrow eN$




Expect 5×10^{10}
 $\tau\tau$ evts...

Qualitative summary





Searches for Charged-Lepton Flavor Violation in Experiments using Intense Muon Beams





Muon g-2

Confirmation of Brookhaven result. Question is now more on Theory...



$$a_\mu(\text{FNAL}) = 116\,592\,040(54) \times 10^{-11} \quad (0.46 \text{ ppm})$$




Flavor physics: quark sector (II)

Lots of attention on Lepton Universality.


A. Greljo @ FAW21*

$$b \rightarrow s \ell^+ \ell^-$$





$$R_K = \mu\mu K / ee K \quad R_{K^*} = \mu\mu K^* / ee K^*$$



+semileptonic + ang. distrib.




B. Capdevila et al @ FAW21*





*Unpaid ad for FAW21: [Flavour Anomaly Workshop 2021](#)

Flavor physics: quark sector (III)

Today: NA62



$$(K^+ \rightarrow \pi^+ \nu \bar{\nu}) = (11.0^{+4.0}_{-3.5}) \times 10^{-11}$$




KOTO 2
KLEVER

NA62
(today+guess)

FCNC in top decays

$BR \times 10^5$	HL-LHC	HE-LHC	ILC ₅₀₀	CLIC ₃₈₀	LHeC	FCC-ee	FCC-hh	FCC-eh
$t \rightarrow Zq$	2.4 - 5.8				4	2.4	≈ 0.1	0.6
$t \rightarrow \gamma c$	7.4		≈ 1	2.6			0.024	
$t \rightarrow \gamma u$	0.86						0.018	
$t \rightarrow \gamma q$					1	1.7		0.085
$t \rightarrow gc$	3.2	0.19						
$t \rightarrow gu$	0.38	0.056						

FCNC with higgs decays



Physics of flavor: Lepton & Quark sectors


□ Neutrinos:

- The very nature of them: Majorana or Dirac?
- Oscillations, (additional) mass generation (beyond EWSB?)
- Richness of the lepton sector (PMNS, CP violation)


□ Quark sector:

- CP violation, CKM triangle
- Rare decays of K and B mesons; SM elements: motivated (universality) and accidental (lepton number conservation)
- Forbidden (?) decays (e.g. $\mu \rightarrow e\gamma$)






Long-lived SUSY?



HL-LHC



Charged stub + photon





DM: summary

- Strengths in WIMP searches both in future lepton and hadron options:
 - Combined FCC program shows best sensitivity to benchmarks
 - Still, needs complementary experiments: DM \neq WIMP (only)
- We can probe the thermal WIMP parameter region
- Large (& yet unknown) parts of phase space can be probed by precision environment/lower bkg in ee



Simplified Models: axial vector

Light DM, $m_\chi = 1\text{ GeV}$



pp: assumes mediator couplings to quarks only.

750 GeV, HL-LHC

1.5 TeV, HE-LHC

3.9 TeV for FCC-hh





Dependence on couplings!

ee: assumes mediator couplings to leptons only. Also in EFT limit, so can be easily rescaled for modified couplings.



Note: taking EFT scale as free parameter, M_{DM} reach \sim kinematic reach of collider.

Significant model dependence. UV models may have comparable quark and lepton couplings. If both present, can also use dilepton resonances.

SUSY: any “holes”?




–: Info only from LHC (& only for stop).
+ : based on data!




Indeed, after LHC, there will be holes [in low mass regions]; closing or looking at how to close them at HL-LHC; for EWKinos, some regions will remain difficult @ pp.

DM: direct detection experiments (I)

Two-phase (liquid) Xe expts continuing to increase their sensitivity




Extending to electron recoils: bkg level: 76×10^{-3} evt/t/yr/keV (1-30 keV)



Tritium contam or Solar Axions ?



Axion explanation? but minuses:
1. very significant model dependence
2. "tension" with astrophysics

Cannot tell, but... sensitivity to 10^{-24} contamination is impressive




L. Hsu ICHEP2020

A. Elykov ICHEP2020 Dark Matter - electron scattering




DM indirect searches: AMS update

The dream scenario




Reality: tantalizing fall (!). But pulsars? Moreover: antiprotons!




Z. Weng ICHEP2020

The future (~2028)



A parting thought: these are (very) difficult measurements; aka: precision makes a difference




Neutrinos

Extras

Neutrinos: Majorana/Dirac ($0\nu\beta\beta$) & mass measurement

As fundamental as a question can get & very hard to answer



Mass limits (measurements...)

Today (BAO + struct. form):
 $\Sigma m_\nu < 0.12$ (95% CL)
 Ultimate (DESI+BAO): $\Sigma m_\nu < 0.02$




Direct m_ν measurement(s)
 KATRIN: $m_\beta < 1.1$ eV (90% CL)
 Asymptotic: $m_\beta < 0.2$ eV
 Ultimate (Project-8): 40 meV

Extremely active field



- Bolometers (CUPID, AMoRE, CANDLES IV)
 - Measure E ($\sigma \sim 0.1$ - 0.3%) from phonons; granularity gives position info
 - Instrumenting with photon detectors for background rejection
- External trackers (SuperNEMO)
 - Trackers + calorimeters, measure E ($\sigma \sim 3$ - 10%) + tracks / positions + PID
- Scintillators (KamLAND2-Zen, SNO+, Theia, ZICOS)
 - Measure E ($\sigma \sim 3$ - 10%) + position from scintillation light; some PID
- Semiconductors (LEGEND, SELENA)
 - Measure E ($\sigma \sim 0.05$ - 0.3%) from ionization; some tracking / position sensitivity
- TPCs (nEXO, NEXT, PandaX, AXEL, NuDEx, DARWIN, LZ)
 - Collect scintillation + ionization: measure E ($\sigma \sim 0.4$ - 3%) + tracks / position + PID

(But) one example: EXO-200 (0.2 t) \rightarrow nEXO (5 t)
 Xe; $T_{1/2} < 3.5 \times 10^{25}$ yr $\rightarrow T_{1/2} < 10^{28}$ yr


(J. Detwiler @ Neutrino2020)



Neutrinos: Mass & nature (Majorana/Dirac – $0\nu\beta\beta$)




Neutrinos: sterile sector (?)



LSND


L~30m
E~30MeV




MiniBooNe
4.6σ ν_e excess!

L~500m
E~500MeV

NEOS



ν_e shortage in
reactor expts
(But TH shaky)




Taking 5 MeV bump
as due to sterile ν

Very active field

Several
expts:
proposed
running

+ICHEP2020
: + DayaBay,
RENO...

Experiment	Reactor Power/Fuel
DANSS (Russia)	3000 MW LEU fuel
NEOS (South Korea)	2800 MW LEU fuel
nulat (USA)	40 MW 235U fuel
Neutrino4 (Russia)	100 MW 235U fuel
PROSPECT (USA)	85 MW 235U fuel
Solid (UK Fr Bel US)	72 MW 235U fuel
Chandler (USA)	72 MW 235U fuel
Stereo (France)	57 MW 235U fuel



M. Danilov
ICHEP 2020

DANSS: excludes RAA
by >8σ (5σ since 2018)

$$P_{\alpha\beta} \simeq \delta_{\alpha\beta} - 4|U_{\alpha\beta}|^2(\delta_{\alpha\beta} - |U_{\alpha\beta}|^2) \sin^2 \left(\frac{\Delta m_{41}^2 L}{4E} \right)$$

$$\nu_e \text{ DIS} \\ \sin^2 2\vartheta_{ee} \simeq 4|U_{e4}|^2$$

$$\nu_\mu \text{ DIS} \\ \sin^2 2\vartheta_{\mu\mu} \simeq 4|U_{\mu 4}|^2$$


$$\nu_\mu \rightarrow \nu_e \text{ APP} \\ \sin^2 2\vartheta_{e\mu} = 4|U_{e4}|^2|U_{\mu 4}|^2 \simeq \frac{1}{4} \sin^2 2\vartheta_{ee} \sin^2 2\vartheta_{\mu\mu}$$

[Okada, Yasuda, IJMPA 12 (1997) 3669; Bilenky, C.G., Grimus, EPJC 1 (1998) 247]


ν_μ → ν_e: LSND, MiniBooNe, OPERA

ν_μ → ν_μ: MiniBooNe, MINOSetal


ν_e → ν_e: Gallium, Reactors, Solar Exp



HL-LHC: Higgs physics



H self-coupling





λ very challenging;
need multiple channels

$$\sigma(m_H) \sim 10\text{-}20 \text{ MeV}$$



Expect $\lesssim 5\%$ uncertainty on main production channels and BRs

Coupling sensitivity expressed as uncertainty on multiplicative modifiers κ_i

$\mathcal{B}(H \rightarrow \text{inv}) = 0.1\%$ in SM, sensitive to BSM, e.g. attractive Dark Matter models



Y_c ?
Not quite



Colliders: the high energy frontier

