



Contribution ID: 66 Contribution code: S01-NPNE-205

Type: Poster presentation

## Determination of $^{226}\text{Ra}$ and $^{222}\text{Rn}$ content in thermomineral water and assessment of radiation risk

*Tuesday, 30 August 2022 18:00 (1h 30m)*

Determination of  $^{226}\text{Ra}$  and  $^{222}\text{Rn}$  activity concentration in water has been recognized as one of the most important tasks in the preservation of public health, since increased content of these radionuclides in drinking water may lead to increased health risk. Generally, the presence of  $^{226}\text{Ra}$  in water samples is potentially dangerous due to its radiotoxicity and has been related to the cancer of bones [1]. On the other hand, its decay product  $^{222}\text{Rn}$  is recognized as a potential cause for the development of lung cancer. Over the decays, several techniques have been developed for the determination of  $^{226}\text{Ra}$  and  $^{222}\text{Rn}$  in water— $\gamma$  spectrometry,  $\alpha$  spectrometry, and liquid scintillation counting.

The concentration of  $^{226}\text{Ra}$  in water depends mostly on the geology of the area and may be elevated if water passes through rocks with elevated content of  $^{238}\text{U}$ . Thus, it is especially important to measure the content of  $^{226}\text{Ra}$  and  $^{222}\text{Rn}$  in thermal and mineral water, since these waters are groundwater and a high concentration of  $^{226}\text{Ra}$  is expected to be found [2]. At the same time, thermal and mineral water is used for different kinds of treatment [3], such as musculoskeletal diseases, arterial hypertension, bone fracture, post-traumatic conditions, neurological disorders, sports injury, etc, which may lead to increased doses due to inhalation and ingestion.

In this work, the activity concentration of  $^{226}\text{Ra}$  and  $^{222}\text{Rn}$  in thermomineral water samples was determined by using the liquid scintillation counting (LSC) technique. The samples were collected from spa Niska Banja [4] located in the southern part of Serbia. A comparison between two methods of sample preparation—one phase and two phases, for  $^{226}\text{Ra}$  determination, was made. At the same time, the  $^{222}\text{Rn}$  content was measured with the active radon device RAD7 and a comparison with the LSC result was conducted. Furthermore, the doses from ingestion and inhalation of  $^{222}\text{Rn}$ , as well as from ingestion of  $^{226}\text{Ra}$  were calculated. The annual effective doses were estimated for patients and tourists and compared with levels for the public imposed by the legislation of the Republic of Serbia.

### References:

1. M. Villa, H.P. Moreno, G. Manjón, Radiat. Meas. 39, 543–550 (2005).
2. J. Mazur et al., Nukleonika 65(2), 77–81 (2020).
3. G. Marović, J. Senčar, Z. Franić, N. Kokobauer, J. Environ. Radioactivity 33(3), 309–317 (1996)
4. J. Nikolov et al., Radiat. Meas. 47, 443–450 (2012).

**Primary authors:** KNEZEVIC RADIC, Jovana (University of Novi Sad, Faculty of Sciences); Dr FORKAPIC, Sofija (University of Novi Sad, Faculty of Sciences); Dr BIKIT, Kristina (University of Novi Sad, Faculty of Sciences); Prof. MRDJA, Dusan (University of Novi Sad, Faculty of Sciences); Mr VELIMIROVIC, Danijel (University of Novi Sad, Faculty of Sciences); Dr HANSMAN, Jan (University of Novi Sad, Faculty of Sciences); Dr KUZMANOVIC, Predrag (University of Novi Sad, Faculty of Sciences); Prof. BIKIT, Istvan (University of Novi Sad, Faculty of Sciences)

**Presenter:** KNEZEVIC RADIC, Jovana (University of Novi Sad, Faculty of Sciences)

**Session Classification:** Poster session

**Track Classification:** Scientific Sections: S01 Nuclear Physics and Nuclear Energy