



Contribution ID: 50 Contribution code: S10-MG-205

Type: Poster presentation (virtual)

## A case study of high PM concentration despite the low anthropogenic pollution in March 2020 during the first COVID-19 lockdown in Sofia

Wednesday, August 31, 2022 11:56 AM (2 minutes)

The aim of this study is to investigate the atmospheric air quality in Sofia during a 2-week period in the beginning of the first COVID-19 lockdown in March 2020. Observations of the aerosol distribution measured with two laser particle counters for PM<sub>2.5</sub> and PM<sub>10</sub> on clear sunny days (7 a.m. to 2 p.m.) at a site inside the built area near the bus stop Pliska on the Tsarigradsko shose during 16-31 March 2020 are presented. Mass and particle number concentrations of PM with high temporal resolution are compared with the data reported by Executive Environmental Agency monitoring network. In the beginning of the period the concentrations of PM<sub>10</sub> vary between 20 and 80  $\mu\text{g}/\text{m}^3$  with a peak around 9 - 10 a.m and PM<sub>2.5</sub> concentrations reach 30  $\mu\text{g}/\text{m}^3$ . On 28 March at 9 a.m. the PM<sub>10</sub> and PM<sub>2.5</sub> concentration raise up to 250  $\mu\text{g}/\text{m}^3$  and 175 ( $\mu\text{g}/\text{m}^3$ ), correspondingly. This abrupt change in the aerosol content in the atmosphere is explained not with anthropogenic load, but with long range dust transport from the Karakum desert region (located between the eastern coast of the Caspian Sea and the steppes near the central Asian mountain ranges), which is a very rare event of intrusion compared to the typical in spring Sahara dust transport. The HYSPLIT back trajectory model is used as additional source of information in the selected days, showing transport in the layer 1000 m – 3000 m from North-West until 26 March and from East on 27-29 March. The GDAS boundary-layer depth at 12 UTC is between 1000 m and 1600 m which suggest intrusion from above during the period. These model data are compared to local meteorological and aerological observations. This study shows the importance of detailed in time laser particle counter observations of PM<sub>10</sub>, PM<sub>2.5</sub> and finer particles for better explanation of causes for the low air quality in urban atmosphere.

**Primary authors:** Prof. BATCHVAROVA, Ekaterina (Climate, Atmosphere and Water Research Institute at Bulgarian Academy of Sciences (CAWRI-BAS)); Dr KOLEV, Nikolay (Department of Physics, University of Mining and Geology “St. Ivan Rilski”, Sofia, Bulgaria); Dr KOLAROVA, Maria (National Institute of Meteorology and Hydrology (NIMH) - BAS, Sofia, Bulgaria); Prof. SAVOV, Plamen (Department of Physics, University of Mining and Geology “St. Ivan Rilski”, Sofia, Bulgaria)

**Presenter:** Dr KOLEV, Nikolay (Department of Physics, University of Mining and Geology “St. Ivan Rilski”, Sofia, Bulgaria)

**Session Classification:** Poster session (virtual)

**Track Classification:** Scientific Sections: S10 Meteorology and Geophysics